

Table des matières

**1. Systematic Reviews and Meta-Analysis** ..... 1

1.1. Generic Acupuncture ..... 1

1.1.1. Sun 2025 ..... 1

1.1.2. Lin 2015 ..... 1

**2. Clinical Practice Guidelines** ..... 2

2.1. American Heart Association/American Stroke Association (AHA, ASA, USA) 2026 Ø ..... 2

# Gait Control After Stroke

## Troubles de la marche post-AVC

### 1. Systematic Reviews and Meta-Analysis

#### 1.1. Generic Acupuncture

##### 1.1.1. Sun 2025

Sun Z, Sun H, Yu K, Zhu Z, Lin Y, Sun K, Zhang J. A Systematic Review and Meta-Analysis of Acupuncture's Impact on Hemiplegic Gait Recovery After Stroke. Complement Ther Med. 2025 Apr 25:103181. <https://doi.org/10.1016/j.ctim.2025.103181>

|            |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                       |
|------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Objective  | This systematic review aims to evaluate the efficacy and safety of acupuncture in alleviating gait disturbances in post-stroke hemiplegia, focusing on various gait parameters.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                       |
| Methods    | A comprehensive search was conducted across PubMed, EMBASE, Cochrane Library, Web of Science, AMED, CINAHL, CBM, CNKI, and WanFang databases to identify relevant randomized controlled trials (RCTs). The included studies were independently evaluated for risk of bias using Cochrane's risk of bias tool. RevMan 5.3 was used for meta-analysis, and adverse events were collected through full-text review.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                      |
| Results    | A total of <b>21 RCTs involving 1463 participants</b> were included. Results showed that acupuncture combined with rehabilitation therapy (RT) significantly improved stride and step length (MD = 7.79, 95% CI: 5.62 to 9.96, Z = 7.03, P < 0.00001, I <sup>2</sup> = 72%), cadence (MD = 10.43, 95% CI: 6.22 to 14.65, Z = 4.85, P < 0.00001, I <sup>2</sup> = 95%), walking speed (MD = 12.27, 95% CI: 9.22 to 15.31, Z = 7.90, P < 0.00001, I <sup>2</sup> = 91%), hip peak flexion angle (MD = 2.71, 95% CI: 0.94 to 4.49, Z = 2.99, P = 0.003, I <sup>2</sup> = 82%), and ankle peak plantarflexion angle (MD = 2.08, 95% CI: 1.11 to 3.06, Z = 4.19, P < 0.0001, I <sup>2</sup> = 0%) compared to RT alone. It also reduced gait cycle time (MD = -0.61, 95% CI: -0.96 to -0.26, Z = 3.44, P = 0.0006, I <sup>2</sup> = 98%) and the proportion of double support phase (MD = -7.16, 95% CI: -9.08 to -5.25, Z = 7.33, P < 0.00001, I <sup>2</sup> = 0%). These improvements in gait parameters suggest enhanced mobility and functional independence for post-stroke patients. However, heterogeneity in participant characteristics and study methodologies was noted, such as variations in stroke types, curses, severity, and acupuncture protocols. The majority of RCTs exhibited moderate to high risk of bias regarding allocation concealment and blinding. Only two RCTs reported no adverse events, while the remaining 19 studies did not mention adverse events. |
| Conclusion | Acupuncture appears to enhance specific aspects of hemiplegic gait, though further high-quality research is needed to fully validate its effects. Current evidence is limited by methodological weaknesses and potential biases in the included studies. Rigorous, well-designed studies are required to further validate the comprehensive effects of acupuncture on post-stroke hemiplegic gait.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                    |

##### 1.1.2. Lin 2015

Lin Meiqin, Liu Weilin. [System review of the efficacy of acupuncture on hemiplegic gait in stroke patients]. Journal of Fujian University of Traditional Chinese Medicine. 2015;1:54-62. [186997].

|                    |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                              |
|--------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Objectives</b>  | In order to obtain objective evidence of acupuncture therapy for stroke patients with hemiplegic gait, we would collect the published studies, and systematic review of the quality, providing reliable evidence for clinical application.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                   |
| <b>Methods</b>     | Computer comprehensively searched Pubmed, EMbase, the Cochrane library, Web of Science, Chinese Science and Technology Database, Wanfang, China National Knowledge Infrastructure, Chinese Biology Medicine and Traditional Chinese Medicine Database. According to bias risk assessment tool of Cochrane, assessed the quality of the studies, evaluated the quality of each study, Re Man 5. 2software was used for meta-analysis and statistical appraisal of effect size.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                |
| <b>Results</b>     | Participants included 22 studies (n=1, 519), <b>8 of randomized controlled trials (RCT)</b> , 14 of controlled clinical trials (CCT). The results suggested that the lower limb Fugl-Meyer assessment scale: at the acute stage which is WMD=6. 71, 95%CI (5. 51, 5. 72) and P<0. 000, 01; at the restoration stage which is WMD=2. 50, 95%CI (1. 49, 3. 50) and P<0. 000, 01; at the sequela stage only 1 trial that has statistical significance (P=0. 003); three trials were aimed at the treatment of post stroke spasm (any course), which is WMD=1. 60, 95%CI (0. 92, 2. 28) and P<0. 000, 01. Acupuncture could improve time-distance parameters, but reported inconsistent results, improve the kinematic parameters of the hip, knee and ankle joint and enhance EMG of back extensor, slow spasm of limbs, increase the balance and walking function, improve the ability of daily life in stroke patients through analysis of quantitative gait. |
| <b>Conclusions</b> | The existing evidences suggested that acupuncture treatment could <b>improve time-distance and kinematic parameters of lower limbs, suppress spasm and increase balance in stroke patients</b> . But the trials involved are low quality, therefore, which remains need to design more rigorous randomized controlled trial to further verification.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                         |

## 2. Clinical Practice Guidelines

⊕ positive recommendation (regardless of the level of evidence reported)  
 ∅ negative recommendation (or lack of evidence)

### 2.1. American Heart Association/American Stroke Association (AHA, ASA, USA) 2026 ∅

Richards LG, Ifejika NL, Stein J, Bahouth MN, Abshire Saylor M, Barrett AM, Chen P, Deutsch AF, Dusenbury W, Ellis C Jr, Eng J, Fleet J, Ford HR, Grattan ES, Harvey RL, Jones CA, Krishnan C, Kutzle S, Matthews D, Mazwi N, Minga J, Nilsen DM, Noser E, Porter HR, Raghavan P, Rose DK, Skidmore ER, St-Onge MP, Stoykov ME, Schettini AT, Terrill AL, Tirschwell DL, Vilar P, Whitaker-Hilbig A, Zorowitz R; Peer Review Committee. 2026 Guideline for Adult Stroke Rehabilitation and Recovery: A Guideline From the American Heart Association and American Stroke Association. Stroke. 2026 Aug 27.

<https://doi.org/10.1161/str.0000000000000536>

- Gait and Mobility: For persons with stroke, **acupuncture** in combination with rehabilitation may be considered for improving gait kinematics and balance. COR 2b; LOE B-R.

From:

<https://ebm.wiki-mtc.org/> - Encyclopédie des sciences médicales chinoises

Permanent link:

<https://ebm.wiki-mtc.org/doku.php?id=acupuncture:evaluation:neuro-psychiatrie:07.%20troubles%20de%20la%20marche%20post-avc> 

Last update: **28 Aug 2026 15:00**